

INFERENCE in the MULTIPLE GAUSSIAN LINEAR MODEL

we will work under the assumption that the model always includes the intercept

$$\underline{x}_1 = \underline{1}_n \text{ with } \beta_1 \text{ the associated coefficient.}$$

1. TEST about an individual coefficient β_j ($j = 2, \dots, p$)

assume that we want to test a single coefficient:

$$\begin{cases} H_0: \beta_j = b_j \\ H_1: \beta_j \neq b_j \end{cases}$$

In particular, we are often interested in testing the statistical significance of an individual coefficient

$$\begin{cases} H_0: \beta_j = 0 \\ H_1: \beta_j \neq 0 \end{cases}$$

- Recall:
- $\hat{\underline{\beta}} \sim N_p(\underline{\beta}, (X^T X)^{-1} \sigma^2)$
 - the j -th element $\hat{\beta}_j \sim N(\beta_j, \underbrace{\sigma^2 [(X^T X)^{-1}]_{jj}}_{V(\hat{\beta}_j)})$
 - $\frac{n \hat{\Sigma}^2}{\sigma^2} \sim \chi^2_{n-p}$
 - $\frac{(n-p) S^2}{\sigma^2} \sim \chi^2_{n-p}$
 - $\hat{\underline{\beta}} \perp \hat{\Sigma}^2 \text{ and } \hat{\underline{\beta}} \perp S^2$

1) We need to define a PIVOTAL QUANTITY

$$\frac{\hat{\beta}_j - b_j}{\sqrt{V(\hat{\beta}_j)}} = \frac{\hat{\beta}_j - b_j}{\sqrt{\sigma^2 [(X^T X)^{-1}]_{jj}}} \stackrel{H_0}{\sim} N(0, 1) \text{ but it depends on the unknown } \sigma^2 \text{ (hence we can't use it)}$$

we consider instead

$$\begin{aligned} T_j &= \frac{\hat{\beta}_j - b_j}{\sqrt{S^2 [(X^T X)^{-1}]_{jj}}} = \frac{\hat{\beta}_j - b_j}{\sqrt{V(\hat{\beta}_j)}} = \\ &= \frac{\hat{\beta}_j - b_j}{\sqrt{\frac{S^2}{\sigma^2} V(\hat{\beta}_j)}} = \frac{\hat{\beta}_j - b_j}{\sqrt{\frac{S^2}{\sigma^2}} \sqrt{\frac{\chi^2_{n-p}}{(n-p)}}} \stackrel{N(0, 1)}{\sim} \\ &\Rightarrow \hat{V}(\hat{\beta}_j) = S^2 [(X^T X)^{-1}]_{jj} \cdot \frac{\sigma^2}{\sigma^2} \\ &= (\sigma^2 [(X^T X)^{-1}]_{jj}) \cdot \frac{S^2}{\sigma^2} = V(\hat{\beta}_j) \cdot \frac{S^2}{\sigma^2} \quad \text{general expression} \end{aligned}$$

$$\Rightarrow T_j = \frac{\hat{\beta}_j - b_j}{\sqrt{V(\hat{\beta}_j)}} \stackrel{H_0}{\sim} t_{n-p}$$

in the simple lm we had $(t-2)$ degrees of freedom. Indeed $p=2$ for the simple lm $X = [\underline{1} \ \underline{x}]$

2) With the data, I compute the OBSERVED VALUE OF THE TEST t_j^{obs}

3) We study the position of the sample space into the REJECT and ACCEPTANCE REGION:

As for the simple linear model, large values of the test (in absolute value) lead to rejecting the null hypothesis (if H_0 is not true, $\hat{\beta}_j$ will be very different from b_j , hence $|\hat{\beta}_j - b_j| \gg 0$ and also $|t^{\text{obs}}| \gg 0$).

Hence: $A_b = (-k, k)$
 $R_b = (-\infty, -k) \cup (k, +\infty)$

4) We conclude the test


4a) FIXED SIGNIFICANCE LEVEL α

$$P_{H_0}(T_j \in R_b) = P_{H_0}(|T_j| > t_{n-p; 1-\frac{\alpha}{2}}) = \alpha$$

i.e.

$$R_b = (-\infty, -t_{n-p; 1-\frac{\alpha}{2}}) \cup (t_{n-p; 1-\frac{\alpha}{2}}, +\infty)$$

and reject H_0 if $|t^{\text{obs}}| \in R_b$



4b) p-value = $P_{H_0}(|T_j| > |t_j^{\text{obs}}|) =$
 $= 2 P_{H_0}(T_j > |t_j^{\text{obs}}|) \text{ with } T_j \sim t_{n-p}$

